Neuroscience Letters, 103 (1989) 354–357
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Advani, A., see Schneiderman, J.H., 39
Aizenman, E., see Rosenberg, P.A., 162
Akaike, N., see Takahashi, K., 29
Akasu, T., see Nishimura, T., 179
Allen, Y.S and Kershaw, T.R., A morphological study of the neuroblastoma-glioma hybrid cell line, NG108-15, in culture and after grafting to the adult rat brain, 247
Angioy, A.M., Tomassini Barbarossa, I., Crnjar, R. and Liscia, A., Effects of octopaminergic substances on the labellar lobe spreading response in the blowfly *Protophormia terraenovae*, 103
Arai, H., Noguchi, I., Sagi, N., Moroji, T. and Iizuka, R., A study of non-isotopic in situ hybridization histochemistry on postmortem changes in vasopressin mRNA in rat brain, 127
Arnold, D., see Schneiderman, J.H., 39
Arora, A., see Millar, T.J., 108

Banati, R.B., see Graeber, M.B., 241
Bargas, J., see Surmeier, D.J., 331
Barnes, C.D., see Fung, S.J., 185
Blakemore, W.F., see Crang, A.J., 269
Bliss, T.B.P., see Lynch, M.A., 191
Bohl, J., see Braak, H., 24
Borgers, M., see Kleiser, B., 225
Braak, E., see Braak, H., 24
Braak, H., Braak, E., Ohm, T. and Bohl, J., Alzheimer's disease: mismatch between amyloid plaques and neuritic plaques, 24
Bradberry, C.W. and Roth, R.H., Cocaine increases extracellular dopamine in rat nucleus acumbens and ventral tegmental area as shown by in vivo microdialysis, 97
Bray, G.A., see Shimizu, H., 74
Bugiani, O., Giaccone, G., Frangione, B., Ghetti, B. and Tagliavini, F., Alzheimer patients: preamyloid deposits are more widely distributed than senile plaques throughout the central nervous system, 263
Bönnemann, C., Holland, A. and Meyer, D.K., Cholecystokinin in rat olfactory bulb is mainly contained in intrinsic neurons, 281

Cammarata, S., see Tabaton, M., 259
Care, J.S., see Wolpaw, J.R., 113
Casamenti, F., Di Patre, P.L., Milan, F., Petrelli, L. and Pepeu, G., Effects of nerve growth factor and GM1 ganglioside on the number and size of cholinergic neurons in rats with unilateral lesion of the nucleus basalis, 87
Chau, R.M.W., see Jen, L.S., 133
Chihara, K., see Jinnai, K., 64
Chorev, M., see Devor, M., 203
Clavelou, P., Pajot, J., Dallel, R. and Raboisson, P., Application of the formalin test to the study of orofacial pain in the rat, 349
Cobcroft, M., Vaccaro, T. and Mitrofanis, J., Distinct patterns of distribution among NADPH-diaphorase neurones of the guinea pig retina, 1
Cools, A.R., see Van den Bos, R., 303
Costa, G., see Polidori, C., 320
Crang, A.J. and Blakemore, W.F., The effect of the number of oligodendrocytes transplanted into X-irradiated glial-free lesions on the extent of oligodendrocyte remyelination, 269
Crnjar, R., see Angioy, A.M., 103

Dallel, R., see Clavelou, P., 349
De Caro, G., see Polidori, C., 320
DeTeresa, R.M., see Masliah, E., 234
Decker, J., see Jorgensen, M.B., 219
Delaunoy, J.P., see Laabich, A., 157
Delay-Goyet, P., Zajac, J.-M. and Roques, B.P., Effects of repeated treatment with haloperidol on rat striatal neutral endopeptidase, and on μ- and δ-opioid binding sites: comparison with chronic morphine and chronic kelatorphan, 197
Devasagayam, T.P.A., Decreased peroxidative potential in rat brain microsomal fractions during ageing, 92
Devor, M., Papir-Kricheli, D., Nachmias, E., Rosenthal, F., Gilon, C., Chorev, M. and Selinger, Z., Substance P-induced cutaneous plasma extravasation in rats is mediated by NK-1 tachykinin receptors, 203
Di Patre, P.L., see Casamenti, F., 87
Dunnett, S.B., see Morris, B.J., 121

Emri, Z., see Juhász, G., 229
Errington, M.L., see Lynch, M.A., 191
Esseling, K., see Kleiser, B., 225

Farber, S.D., see Low, W.C., 314
Fastbom, J., see Rudolphi, K.A., 275
Fluri, M., see Ruegg, D.G., 174
Frangione, B., see Bugiani, O., 263

0304-3940/89/$ 03.50 © 1989 Elsevier Scientific Publishers Ireland Ltd.

Fredholm, B.B., see Rudolphi, K.A., 275
Fujita, T., see Jinnai, K., 64
Fung, S.J. and Barnes, C.D., Raphé-produced excitation of spinal cord motoneurons in the cat, 185

Geppetti, P., Santicioli, P., Rubini, I., Spillantini, M.G., Maggi, C.A. and Sicuteri, F., Thiorphan increases capsaicin-evoked release of substance P from slices of dorsal spinal cord of guinea pig, 69
Ghetti, B., see Bugiani, O., 263
Giaccone, G., see Bugiani, O., 263
Gilon, C., see Devor, M., 203
Graeber, M.B., Banati, R.B., Streit, W.J. and Kreutzberg, G.W., Immunophenotypic characterization of rat brain macrophages in culture, 241
Gregory, J.E., McIntyre, A.K. and Proske, U., Tendon organ afferents in the knee joint nerve of the cat, 287
Guerrero, J.M., see Santana, C., 338

Hajós, F., see Kardos, J., 139
Hansen, L.A., see Masliah, E., 234
Haseldonckx, M., see Kleiser, B., 225
Heidenreich, B.A. and Rebec, G.V., Serotonergic dorsal raphe neurons: changes in spontaneous neuronal activity and responsiveness to 5-MeODMT following long-term amphetamine administration, 81
Hernandez, D.E. and Xue, B.G., Imipramine prevents stress gastric glandular lesions in rats, 209
Hill, T.G., see Low, W.C., 314
Holland, A., see Bönnemann, C., 281
Horcholle-Bossavit, G., Jami, L., Peptit, J., Vejsada, R. and Zytnicki, D., Activation of Golgi tendon organs by asynchronous contractions of motor units in cat leg muscles, 44
Horn, E., see Kleiser, B., 225
Houle, J.D. and Johnson, J.E., Nerve growth factor (NGF)-treated nitrocellulose enhances and directs the regeneration of adult rat dorsal root axons through intraspinal neural tissue transplants, 17
Houle, J.D. and Reier, P.J., Regrowth of calcitonin gene-related peptide (CGRP) immunoreactive axons from the chronically injured rat spinal cord into fetal spinal cord tissue transplants, 253
Hughes, S. and Smith, M.E., Proopiomelanocortin-derived peptides in mice with motoneurone disease, 169

Iizuka, R., see Arai, H., 127
Imamura, Y., see Ishikawa, K., 343
Ishikawa, K., Shibanoki, S., Kubo, T., Kogure, M., Imamura, Y., Osawa, N., Ohmura, M. and Mikoshiba, K., Functional difference in monoamine transmitters in the behaviourally abnormal mouse mutant (*wriggle mouse sagami*), 343

Jami, L., see Horcholle-Bossavit, G., 44
Jen, L.S., Zhao, L.P. and Chau, R.M.W., Cytochrome oxidase activity in the rat retina following unilateral thalamic lesion, 133
Jinnai, K., Chihara, K., Kanda, F., Tada, K. and Fujita, T., Calcitonin gene-related peptide enhances spontaneous acetylcholine release from the rat motor nerve terminal, 64
Johnson, J.E., see Houle, J.D., 17
Jorgensen, M.B., Decker, J. and Wright, D.C., Binding of [^{3}H]inositoltrisphosphate and [^{3}H]phorbol 12,13-dibutyrate in rat hippocampus following transient global ischemia: a quantitative autoradiographic study, 219
Juhász, G., Emri, Z. and Kékesi, K., Local perfusion of the thalamus with GABA increases sleep and induces long-lasting inhibition of somatosensory event-related potentials in cats, 229

Kamiya, H., see Swada, S., 34
Kanda, F., see Jinnai, K., 64
Kaneda, M., see Takahashi, K., 29
Kardos, J., Kovács, I., Hajós, F., Kálmán, M. and Simonyi, M., Nerve endings from rat brain tissue release copper upon depolarization. A possible role in regulating neuronal excitability, 139
Kawahara, K., Nakazono, Y., Yamauchi, Y. and Miyamoto, Y., Coupling between respiratory and locomotor rythms during fictive locomotion in decerebrate cats, 326
Keil, M., see Rudolphi, K.A., 275
Kershaw, T.R., see Allen, Y.S., 247
Kitai, S.T., see Surmeier, D.J., 331
Kitao, Y., see Nakamura, Y., 145
Kleiser, B., Van Reempts, J., Van Deuren, B., Haseldonckx, M., Borgers, M., Horn, E., Esseling, K., Widder, B. and Kornhuber, H.J., Favourable effect of flunarizine on the recovery from hemiparpesis in rats with intracerebral hematomas, 225
Kogure, M., see Ishikawa, K., 343
Koob, G.F., see Pulvirenti, L., 213

Kornhuber, H.J., see Kleiser, B., 225
Kovács, I., see Kardos, J., 139
Kreutzberg, G.W., see Graeber, M.B., 241
Kubek, M.J., see Low, W.C., 314
Kubo, T., see Ishikawa, K., 343
Kudo, M., see Nakamura, Y., 145
Kuo, C.-H. and Miki, N., Translocation of a photoreceptor-specific MEKA protein by light, 8
Kálmán, M., see Kardos, J., 139
Kékesi, K., see Juhász, G., 229

Laabich, A., Sensenbrenner, M. and Delaunoy, J.P., Monolayer cultures of ependymal cells on orous bottom dishes. A tool for transport studies across the brain cerebrospinal barrier, 157
Leanza, Z., Pellitteri, R., Russo, A. and Stanzani, S., Branching projections from subcoeruleus area neurons to medial preoptic area and cervical spinal cord revealed by double retrograde neuronal labeling, 11
Lee, C.L., see Wolpaw, J.R., 113
Liscia, A., see Angioy, A.M., 103
Low, W.C., Roepke, J., Farber, S.D., Hill, T.G., Sattin, A. and Kubek, M.J., Distribution of thyrotropin-releasing hormone (TRH) in the hippocampal formation as determined by redioimmunoassay, 314
Lynch, M.A., Errington, M.L. and Bliss, T.B.P., The increase in [^{3}H]glutamate release associated with long-term potentiation in the dentate gyrus is blocked by commissural stimulation, 191

Maggi, C.A., see Geppetti, P., 69
Mancardi, G., see Tabaton, M., 259
Manetto, V., see Tabaton, M., 259
Masliah, E., Terry, R.D., DeTeresa, R.M. and Hansen, L.A., Immunohistochemical quantification of the synapse-related protein synaptophysin in Alzheimer disease, 234
Massi, M., see Polidori, C., 320
McIntyre, A.K., see Gregory, J.E., 287
Meyer, D.K., see Bönnemann, C., 281
Miki, N., see Kuo, C.-H., 8
Mikoshiba, K., see Ishikawa, K., 343
Milan, F., see Casamenti, F., 87
Millar, T.J., Vaegan, and Arora, A., Urethane as a sole general anaesthetic in cats used for electroretinogram studies, 108
Millar, T.J., see Yang, G., 151
Mitrofanis, J., see Cobcroft, M., 1
Miyamoto, Y., see Kawahara, K., 326
Morgan, I.G., see Yang, G., 151
Moriizumi, T., see Nakamura, Y., 145
Moroji, Y., see Arai, H., 127
Morris, B.J., Wisden, W., Dunnett, S.B. and Sirinathsinghji, D.J.S., Cellular localisation of somatostatin mRNA and neuropeptide Y mRNA in foetal striatal tissue grafts, 121
Murase, K., Ryu, P.D. and Randic, M., Excitatory and inhibitory amino acids and peptide-induced responses in acutely isolated rat spinal dorsal horn neurons, 56

Nachmias, E., see Devor, M., 203
Nagatsu, I., see Nishimura, T., 179
Nakamura, Y., Tokuno, H., Moriizumi, T., Kitao, Y. and Kudo, M., Monosynaptic nigral inputs to the pedunculopontine tegmental nucleus neurons which send their axons to the medial reticular formation in the medulla oblongata. An electron microscopic study in the cat, 145
Nakazono, Y., see Kawahara, K., 326
Nishimura, T., Yoshida, M., Nagatsu, I. and Akasu, T., Frequency dependent inhibition of the nicotinic transmission by serotonin in vesical pelvic ganglia of the rabbit, 179
Noguchi, I., see Arai, H., 127

Ohm, T., see Braak, H., 24
Ohmura, M., see Ishikawa, K., 343
Osawa, N., see Ishikawa, K., 343

Pagni, S., see Romaiguère, P., 50
Pajot, J., see Clavelou, P., 349
Papir-Kricheli, D., see Devor, M., 203
Pellitteri, R., see Leanza, Z., 11
Pepeu, G., see Casamenti, F., 87
Peptit, J., see Horcholle-Bossavit, G., 44
Perfumi, M., see Polidori, C., 320
Perry, G., see Tabaton, M., 259
Petrelli, L., see Casamenti, F., 87
Polidori, C., Saija, A., Perfumi, M., Costa, G., De Caro, G. and Massi, M., Vasopressin release induced by intracranial injection of tachykinins is due to activation of central neurokinin-3 receptors, 320
Prasad, C. and Spahn, S.A., Chronic thyrotropin-releasing hormone decreases the affinity and increases the number of its own receptor in the spinal cord, 309
Proske, U., see Gregory, J.E., 287
Pulvirenti, L., Swerdlow, N.R. and Koob, G.F., Microinjection of a glutamate antagonist into the nucleus accumbens reduces psychos + K1 ce-

rebral cortex, 162

Rosenthal, F., see Devor, M., 203
Roth, R.H., see Bradberry, C.W., 97
Rubini, I., see Geppetti, P., 69
Rudolphi, K.A., Keil, M., Fastbom, J. and Fredholm, B.B., Ischaemic damage in gerbil hippocampus is reduced following upregulation of adenosine (A_1) receptors by caffeine treatment, 275
Ruegg, D.G. and Fluri, M., Relation between electromyogram and torque of isometric reflex contractions in man, 174
Ruppersberg, J.P., see Siara, J., 298
Russo, A., see Leanza, Z., 11
Ryu, P.D., see Murase, K., 56
Rüdel, R., see Siara, J., 298

Sagi, N., see Arai, H., 127
Saija, A., see Polidori, C., 320
Santana, C., Guerrero, J.M. and Reiter, R.J., Effects of either forskolin, the 1,9-dideoxy derivative of forskolin, or 8-bromocyclic AMP on cyclic AMP and melatonin production in the Syrian hamster pineal gland in organ culture, 338
Santicioli, P., see Geppetti, P., 69
Sattin, A., see Low, W.C., 314
Schneiderman, J.H., Arnold, D. and Advani, A., Power spectral analysis of spontaneous field potentials in hipocampal slice, 39
Selinger, Z., see Devor, M., 203
Sensenbrenner, M., see Laabich, A., 157
Shibanoki, S., see Ishikawa, K., 343
Shimizu, H. and Bray, G.A., Modulation by corticotropin-releasing factor of monoamine metabolism in the lateral hypothalamus, 74
Siara, J., Ruppersberg, J.P. and Rüdel, R., Human acetylcholine receptors desensitize much faster than rat acetylcholine receptors, 298
Sicuteri, F., see Geppetti, P., 69
Simonyi, M., see Kardos, J., 139
Sirinathsinghji, D.J.S., see Morris, B.J., 121
Smith, M.E., see Hughes, S., 169
Spahn, S.A., see Prasad, C., 309
Spillantini, M.G., see Geppetti, P., 69
Stanzani, S., see Leanza, Z., 11
Streit, W.J., see Graeber, M.B., 241
Surmeier, D.J., Bargas, J. and Kitai, S.T., Two types of A-current differing in voltage-dependent are expressed by neurons of the rat neostriatum, 331
Swada, S., Kamiya, H. and Yamamoto, C., Simultaneous recording of presynaptic spikes and excitatory postsynaptic potentials from monosynaptically connected hippocampal neurons, 34
Swerdlow, N.R., see Pulvirenti, L., 213

Tabaton, M., Cammarata, S., Manetto, V., Perry, G. and Mancardi, G., Tau-reactive neurofibrillary tangles in cerebellar cortex from patients with Alzheimer's disease, 259
Tada, K., see Jinnai, K., 64
Tagliavini, F., see Bugiani, O., 263
Takahashi, K., Tateishi, N., Kaneda, M. and Akaike, N., Comparison of low-threshold Ca^{2+} currents in the hippocampal CA1 neurons among the newborn, adult and aged rats, 29
Tateishi, N., see Takahashi, K., 29
Terry, R.D., see Masliah, E., 234
Tokuno, H., see Nakamura, Y., 145
Tomassini Barbarossa, I., see Angioy, A.M., 103

Vaccaro, T., see Cobcroft, M., 1
Vaegan, see Millar, T.J., 108
Van Deuren, B., see Kleiser, B., 225
Van Reempts, J., see Kleiser, B., 225
Van den Bos, R. and Cools, A.R., Involvement of the substantia innominata/ventral pallidum complex in transmitting forelimb muscular rigidity evoked from the nucleus accumbens in rats, 303
Vedel, J.-P., see Romaiguère, P., 50
Vejsada, R., see Horcholle-Bossavit, G., 44
Vyskocil, F., see Zemková, H., 293

Widder, B., see Kleiser, B., 225
Wisden, W., see Morris, B.J., 121
Wolpaw, J.R., Care, J.S. and Lee, C.L., Memory traces in spinal cord produced by H-reflex conditioning: effects of post-tetanic potentiation, 113
Wright, D.C., see Jorgensen, M.B., 219